



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2008

The anatomy of Leo I: how tidal tails affect the kinematics

Lokas, E L ; Klimentowski, J ; Kazantzidis, S ; Mayer, L

Abstract: We model the recently published kinematic data set for Leo I dwarf spheroidal (dSph) galaxy by fitting the solutions of the Jeans equations to the velocity dispersion and kurtosis profiles measured from the data. We demonstrate that when the sample is cleaned of interlopers the data are consistent with the assumption that mass follows light and isotropic stellar orbits with no need for an extended dark matter halo. Our interloper removal scheme does not clean the data of contamination completely, as demonstrated by the rotation curve of Leo I. When moving away from the centre of the dwarf, the rotation appears to be reversed. We interpret this behaviour using the results of an N-body simulation of a dwarf galaxy possessing some intrinsic rotation, orbiting in the Milky Way potential and show that it can be reproduced if the galaxy is viewed almost along the tidal tails so that the leading (background) tail contaminates the western part of Leo I while the trailing (foreground) tail the eastern one. We show that this configuration leads to a symmetric and Gaussian distribution of line-of-sight velocities. The simulation is also applied to test our modelling method on mock data sets. We demonstrate that when the data are cleaned of interlopers and the fourth velocity moment is used the true parameters of the dwarf are typically within 1 errors of the best-fitting parameters. Restricting the fitting to the inner part of Leo I our best estimate for the anisotropy is $-0.2^{+0.3}_{-0.4}$ and the total mass $M = (4.5 \pm 0.7) \times 10^7 M_{\odot}$. The mass-to-light ratio (M/L) including the errors in mass, brightness and distance is $M/L = 8.2 \pm 4.5$ solar units.

DOI: <https://doi.org/10.1111/j.1365-2966.2008.13661.x>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-16542>

Journal Article

Accepted Version

Originally published at:

Lokas, E L; Klimentowski, J; Kazantzidis, S; Mayer, L (2008). The anatomy of Leo I: how tidal tails affect the kinematics. *Monthly Notices of the Royal Astronomical Society*, 390(2):625-634.

DOI: <https://doi.org/10.1111/j.1365-2966.2008.13661.x>

The anatomy of Leo I: how tidal tails affect the kinematics

Ewa L. Łokas,¹ Jarosław Klimontowski,¹ Stelios Kazantzidis² and Lucio Mayer^{3,4}

¹*Nicolaus Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland*

²*Center for Cosmology and Astro-Particle Physics; and Department of Physics; and Department of Astronomy, The Ohio State University, Physics Research Building, 191 West Woodruff Avenue, Columbus, OH 43210, USA*

³*Institute for Theoretical Physics, University of Zürich, CH-8057 Zürich, Switzerland*

⁴*Institute of Astronomy, Department of Physics, ETH Zürich, Wolfgang-Pauli Strasse, CH-8093 Zürich, Switzerland*

3 September 2010

ABSTRACT

We model the recently published kinematic data set for Leo I dSph galaxy by fitting the solutions of the Jeans equations to the velocity dispersion and kurtosis profiles measured from the data. We demonstrate that when the sample is cleaned of interlopers the data are consistent with the assumption that mass follows light and isotropic stellar orbits with no need for an extended dark matter halo. Our interloper removal scheme does not clean the data of contamination completely, as demonstrated by the rotation curve of Leo I. When moving away from the centre of the dwarf, the rotation appears to be reversed. We interpret this behaviour using the results of an N -body simulation of a dwarf galaxy possessing some intrinsic rotation, orbiting in the Milky Way potential and show that it can be reproduced if the galaxy is viewed almost along the tidal tails so that the leading (background) tail contaminates the western part of Leo I while the trailing (foreground) tail the eastern one. We show that this configuration leads to a symmetric and Gaussian distribution of line-of-sight velocities. The simulation is also applied to test our modelling method on mock data sets. We demonstrate that when the data are cleaned of interlopers and the fourth velocity moment is used the true parameters of the dwarf are typically within 1σ errors of the best-fitting parameters. Restricting the fitting to the inner part of Leo I our best estimate for the anisotropy is $\beta = -0.3^{+0.4}_{-0.6}$ and the total mass $M = (3.6 \pm 0.6) \times 10^7 M_\odot$. The mass-to-light ratio including the errors in mass, brightness and distance is $M/L_V = 6.6 \pm 3.7$ solar units.

Key words: galaxies: Local Group – galaxies: dwarf – galaxies: clusters: individual: Leo I – galaxies: fundamental parameters – galaxies: kinematics and dynamics – cosmology: dark matter

1 INTRODUCTION

The Leo I dwarf spheroidal (dSph) galaxy discovered by Harrington & Wilson (1950) is one of the brightest and most distant members of the Local Group dSph galaxy population. Its large distance and significant velocity directed away from the Milky Way make its dynamical status still unclear, both in terms of whether it is bound to either the Milky Way or M31 (Byrd et al. 1994) and to what extent its internal dynamics may be affected by tidal interactions with the host galaxy.

The heliocentric velocity of Leo I was first determined from a single carbon star by Aaronson, Hodge & Olszewski (1983) which was later followed by measurements for red giants by Suntzeff et al. (1986). Mateo et al. (1998) estimated with good accuracy radial velocities for 33 red giants which allowed them to determine for the first time the galaxy's

velocity dispersion and mass-to-light ratio (M/L) of 3.5–5.6 solar units in V -band. Although lower than in other dSph galaxies, this value of M/L indicates the presence of a significant amount of dark matter given the relatively young stellar population of Leo I (Lee et al. 1993; Caputo et al. 1999; Gallart et al 1999).

During the past year, three studies on Leo I dynamics have appeared in the literature: Koch et al. (2007, hereafter K07), Sohn et al. (2007, hereafter S07) and Mateo, Olszewski & Walker (2008, hereafter M08). Each presented new kinematic measurements for Leo I stars and discussed their interpretation. No consistent image of the galaxy dynamics however emerged from these studies and their conclusions were on many points contradictory. First, while S07 and M08 estimated the mass-to-light ratio of 10 solar units in the V -band or lower, K07 found a value as high as 24.




Figure 1. Upper panels: the kinematic samples of Leo I stars used in the modelling. The left panel shows the original sample of 328 stars identified as members by M08, while the right one presents the sample of 316 stars obtained by rejection of interlopers (open circles). The middle (lower) panels show the velocity dispersion (kurtosis) profiles obtained from the corresponding samples with binning $9 \times 30 + 2 \times 29$ for the sample of 328 stars and $4 \times 31 + 6 \times 32$ for the sample of 316 stars. The dashed lines show the best-fitting dispersion profiles when only the dispersion is fitted, while the solid lines plot the best-fitting profiles of the moments when the dispersion and kurtosis are fitted simultaneously. The thinner lines were obtained from fitting all data points, the thicker ones with the three outer data points rejected.

In addition, M08 and K07 claimed that the kinematic data are inconsistent with a simple hypothesis that mass follows light and require an extended dark matter halo. Second, all three investigations reported the detection of rotation at some level, but different interpretations of this finding were given by S07 and M08: M08 concluded that the western part of Leo I showing stellar velocities positive with respect to the mean is affected by the leading tidal tail and the eastern part with velocities below the mean by the trailing tail; the interpretation of S07 placed the corresponding tails in opposite directions. Third, while S07 found the velocity distribution of their kinematic sample to be asymmetric and interpreted it as a signature of tidal interaction, the distribution of the stellar sample of M08 is symmetric and Gaussian-like. The purpose of this work is to explain the differences and propose

a detailed model for the origin of the kinematic properties of Leo I.

The paper is organized as follows. In section 2 we present detailed models of Leo I kinematics assuming that mass follows light and using the data set of M08. The data are modelled by fitting the velocity dispersion and kurtosis profiles. We also discuss the possible contamination of the data by stars from the tidal tails. We demonstrate that this contamination manifests itself not only in the overestimated values of the outer dispersion data points but also in the shape of the galaxy’s rotation curve. In section 3 we use a collisionless N -body simulation of a dwarf galaxy orbiting in the Milky Way potential to propose a detailed model of Leo I including the orientation of its tidal tails with respect to the observer and show that such a configuration leads to a symmetric velocity distribution. We also use the simulation

Table 1. Adopted parameters of Leo I.

parameter	value
centre	RA=10 ^h 08 ^m 27 ^s Dec=+12°18′30″
distance modulus $(m - M)_0$	22.02 ± 0.13
distance D	254 ± 15 kpc
apparent magnitude m_V	10.0 ± 0.3
absolute magnitude M_V	-12.02 ± 0.43
luminosity L_V	$(5.5 \pm 2.2) \times 10^6 L_\odot$
Sérsic radius R_S	2.0 arcmin
Sérsic parameter m	1.0
major axis PA	79°

to generate mock data sets and model them in order to verify the reliability of our method. The discussion, including a detailed comparison with earlier work, follows in section 4.

2 MODELLING OF LEO I

Figure 1 shows in the upper left panel the kinematic sample of 328 stars from M08. The diagram plots the heliocentric velocities of Leo I stars as a function of distance from the centre of the galaxy which we also adopt from M08 (see Table 1). The selection of these 328 stars out of the entire sample of 387 stars listed in table 5 of M08 was done by rejecting obvious outliers including pronounced contribution from the Milky Way stars (see below). From these data we calculated the velocity dispersion profile $\sigma(R)$ shown in the middle left panel of Fig. 1 in 11 radial bins of $9 \times 30 + 2 \times 29$ stars using a standard unbiased estimator of dispersion (see e.g. Lokas, Mamon & Prada 2005). The data points were assigned sampling errors of size $\sigma/\sqrt{2(n-1)}$ where n is the number of stars per bin. The lower left panel plots the kurtosis-like variable $k = (\log \kappa)^{1/10}$ which has a Gaussian sampling distribution contrary to the kurtosis κ . The values of k were obtained with the correction of the standard estimator of kurtosis K by the bias due to the low number of stars per bin so that $\kappa = 3K/2.68$. The data points were assigned sampling errors of 2 percent (see Lokas & Mamon 2003; Lokas et al. 2005).

For the modelling we made the simplest possible assumption that mass follows light or equivalently that M/L is constant with radius. The light distribution in terms of the Sérsic (1968) profile (for the formulae see Lokas et al. 2005) with $R_S = 2.0$ arcmin and $m = 1$ (which is equivalent to the exponential profile) was adopted from Irwin & Hatzidimitriou (1995). We also adopt their total apparent magnitude which we translate into the absolute value using the distance of Leo I $D = 254$ kpc as estimated by Bellazzini et al. (2004) from the tip of the red giant branch (which agrees well with the distance found by Held et al. 2001 from the RR Lyrae stars). The values of all the adopted parameters with errors are listed in Table 1. The error in luminosity includes the error in the measured apparent magnitude as well as the distance.

We modelled the velocity moments using the solutions of the Jeans equations as described in Lokas (2002) and Lokas et al. (2005) adjusting two free parameters, the total mass and the anisotropy parameter β which was as-





Figure 2. The 1σ , 2σ and 3σ confidence regions in the $M - \beta$ parameter plane obtained from fitting the dispersion (left panels) and both dispersion and kurtosis (right panels). The text in each panel specifies the sample for which the fit was performed. The best-fitting parameters (marked with dots) are listed in Table 2 with 1σ error bars.

sumed to be constant with radius. The best-fitting solutions in the case when only the dispersion profile is considered are plotted as dashed lines in Fig. 1 and the corresponding confidence regions in the $M - \beta$ parameter plane following from the sampling errors are illustrated in the left column of Fig. 2. The solid lines in Fig. 1 show the resulting best-fitting profiles in the case when both dispersion and kurtosis profiles are fitted simultaneously. The corresponding confidence contours are plotted in the right column of Fig. 2. For all cases the best-fitting parameters with 1σ error bars are listed in Table 2.

The contours shown in the upper left panel of Fig. 2

Table 2. Fitted parameters of Leo I.

sample	fitted	β	$M[10^7 M_\odot]$	$M/L_V[M_\odot/L_\odot]$	χ^2/N
328 stars, 11 data points	σ	$-2.4^{+1.4}_{-5.6}$	4.2 ± 0.6	7.7 ± 4.2	21.0/9
	$\sigma + \kappa$	$-0.9^{+0.5}_{-0.8}$	4.4 ± 0.6	8.0 ± 4.3	35.4/20
328 stars, 8 data points	σ	$-0.5^{+0.5}_{-0.9}$	3.9 ± 0.6	7.0 ± 3.9	3.2/6
	$\sigma + \kappa$	$-0.4^{+0.5}_{-0.6}$	3.9 ± 0.6	7.1 ± 3.9	10.6/14
316 stars, 10 data points	σ	$-1.1^{+0.7}_{-1.7}$	3.9 ± 0.6	7.1 ± 3.9	13.3/8
	$\sigma + \kappa$	$-0.8^{+0.5}_{-0.7}$	4.0 ± 0.5	7.3 ± 3.8	17.4/18
316 stars, 7 data points	σ	$-0.2^{+0.4}_{-0.7}$	3.6 ± 0.6	6.6 ± 3.7	1.5/5
	$\sigma + \kappa$	$-0.3^{+0.4}_{-0.6}$	3.6 ± 0.6	6.6 ± 3.7	4.2/12

**Figure 3.** Rotation curves of Leo I stars along the major axis of the photometric image. The left panel shows the original sample of 328 stars identified as members by M08, while the right one presents the sample of 316 stars obtained by the interloper removal method of den Hartog & Katgert (1996). The binning is $9 \times 30 + 2 \times 29$ for the sample of 328 stars and $4 \times 31 + 6 \times 32$ for the sample of 316 stars (from negative to positive X). The horizontal line in each panel marks the mean velocity of each sample.

correspond to the result obtained by M08, namely that if we force the assumption that mass follows light then the inferred anisotropy will be very tangential. In addition, the overall fit is quite bad (with $\chi^2/N = 21.0/9$, see Table 2). This is obviously caused by the larger values of the outer 3 dispersion data points. Interestingly, when the kurtosis is added to the analysis, the best-fitting anisotropy is much less tangential (see the upper right panel of Fig. 2) but the quality of the fit is still bad ($\chi^2/N = 35.4/20$). The secondary increase of the velocity dispersion profile at larger projected radii R may be interpreted as a signature of an extended dark matter halo but also as due to the contamination by stars from the tidal tails if they are aligned with the observer's line of sight.

As discussed in detail by Klimontowski et al. (2007), these stars will contribute significantly to artificially inflate the velocity dispersion mainly in the outer radial bins. If this is the case, it is advisable (as suggested by M08 themselves) to use only the inner part of the velocity dispersion profile. We have therefore repeated the analysis rejecting the outer three bins in the velocity dispersion and kurtosis profiles. The best-fitting solutions of the Jeans equations for such truncated moments are plotted with thicker lines in Fig. 1 ending at the last point included in the fit. Again the dashed and solid lines correspond to the fits done for the dispersion alone and for both moments. Note that the

inferred anisotropy is now (second row of panels in Fig. 2) consistent with isotropy at 1σ level, and the kurtosis only helps to constrain the anisotropy more strongly.

Klimontowski et al. (2007) demonstrated that the contamination from the tidal tails can be at least partially removed from the kinematic data sets by adopting the interloper removal method of den Hartog & Katgert (1996) originally devised for galaxy clusters. The method turned out to work very effectively on mock kinematic data sets generated from a simulated dwarf galaxy being tidally stripped by the Milky Way potential removing most of unbound stars from the tidal tails present in the data due to projection effects. Applying this method to the present sample for Leo I we remove 12 stars marked in the upper right panel of Fig. 1 as open circles. The corresponding velocity moments calculated for this reduced sample of 316 stars are also shown in the right column of the Figure. We repeated the fitting of the moments as for the entire sample and the results are shown in Figs. 1 and 2 as well as Table 2 in an analogous way. We can see that the dispersion in the outer bins is now significantly reduced and even when all 10 data points are fitted the quality of the fit is acceptable (only the outermost dispersion point with $R > 10$ arcmin is really discrepant). Note also that the best-fitting masses are now somewhat lower than for the sample of 328 stars.

When the fit is restricted to the inner 7 data points of

each velocity moment the quality is further improved and the results are fully consistent with isotropy of stellar orbits and the hypothesis of mass following light. As the final results of our analysis we suggest adopting those obtained for the most reliable sample of 7 inner data points for each moment calculated from the set of 316 stars. From fitting both velocity moments we obtain in this case with $\chi^2/N = 4.2/12$ the anisotropy $\beta = -0.3^{+0.4}_{-0.6}$ and the total mass $M = (3.6 \pm 0.6) \times 10^7 M_\odot$. The quoted errors are the 1σ errors following from the sampling errors of velocity moments. Combining this mass with the luminosity from Table 1 we get the mass-to-light ratio $M/L_V = 6.6 \pm 3.7 M_\odot/L_\odot$ where the error includes the error in mass, the measurement of the apparent magnitude and the distance (the values of M/L_V for other fitted cases are listed in Table 2). Since the stellar mass-to-light ratio of the relatively young stellar population of Leo I is estimated to be below $1 M_\odot/L_\odot$ this value points to the presence of a significant amount of dark matter. We conclude however that the kinematic data for Leo I can be explained without an extended dark matter halo as the assumption of mass following light works quite well.

In order to verify our hypothesis that the kinematic data set for Leo I is indeed contaminated by stars from the tidal tails we propose to consider the rotation curve obtained from the same data. The curves are obtained by binning the velocities in a similar way as before but along the major axis of the dwarf (assumed to lie at PA = 79° , as determined by Irwin & Hatzidimitriou 1995). The results are shown in Fig. 3 again for the total sample of 328 stars and the cleaned sample with 316 stars. As we can see, the inner parts of the diagrams ($|X| < 3$ arcmin) are consistent with weak rotation such that the western part is approaching and the eastern receding. The direction of the rotation is reversed however when we go farther out from the centre of the dwarf. Interestingly, this result agrees well with what was reported in S07 (see their fig. 17) despite the fact that their data cover only a fraction of the dwarf area on the sky (two differently oriented rectangles on the eastern and western side of the dwarf, see their fig. 16). The rotation curve shown in Fig. 3 also agrees with that of the eastern part shown in fig. 3 of K07 (in the western part their data show no rotation).

3 COMPARISON WITH *N*-BODY SIMULATIONS

3.1 Rotation and tidal tails

In order to verify whether such a reversed rotation can be due to tidal interactions we have resorted to an *N*-body simulation. We used the last output of the simulation described in Klimentowski et al. (2007). The simulation followed the evolution of a two-component (stars and dark matter) dwarf orbiting in the Milky Way potential. Over 10 Gyr of evolution the dwarf completes five eccentric orbits losing ~ 99 percent of its initial mass. The tidal interactions lead to the formation of pronounced tidal tails which are present for most of the time. In the final output, the dwarf is at the apocentre, its shape is spheroidal, and the tidal tails are oriented approximately towards the centre of the Milky Way (Klimentowski et al. 2008).




Figure 4. Upper left: the simulated dwarf as seen on the sky. Upper right: the view of the dwarf from above. The observer sees the galaxy along the y axis ($x = 0$) from below (more negative y). Lower panel: the rotation curve measured by the observer from stars seen within $|x| < 3$ kpc, $|z| < 2.5$ kpc and with velocities within ± 25 km s $^{-1}$ with respect to the dwarf's mean. The solid line shows the curve obtained with all stars, the dashed one from the combined 100 samples of 200 stars after application of the interloper removal scheme. The dotted line shows the rotation curve not affected by tidal tails, obtained by selecting stars within $|y| < 2.5$ kpc.

The orbital apocentre of the simulated dwarf is 120 kpc which corresponds to a distance that is twice smaller compared to the current distance of Leo I. Note however that a typical cosmological orbit for satellites with apocentre to pericentre ratio $r_{\text{apo}}/r_{\text{peri}} \sim 6/1$ and r_{apo} comparable to the current distance of Leo I should still allow the transformation from a disk to a spheroid to be completed after about 10 Gyr (Mayer et al. 2001). The current orbit of Leo I might also be the result of scattering from an orbit with much higher binding energy where the transformation might have been much more efficient with the original apocentre much smaller (Sales et al. 2007; M08).

Given that dwarfs spend most of their orbital time near apocentre, the state of the simulated dwarf should be *qualitatively* similar to that of Leo I. An observer situated near the Milky Way will view the dwarf nearly along its tidal tails (Klimentowski et al. 2008). The configuration is illustrated in Fig. 4 with the upper left panel showing the dwarf as it appears to the observer on the sky and the upper right panel showing the view from above the dwarf so that the line of sight is along the axis $x = 0$ and the observer is looking from the direction of negative y . From such a configuration we choose stars with $|x| < 3$ kpc and $|z| < 2.5$ kpc and project their velocities along the y axis to produce line-of-sight velocities available for observation. We also introduce a cut-off in these velocities at ± 25 km s $^{-1}$ with respect to the mean velocity of the dwarf, which corresponds to $\pm 4\sigma$ range in velocities where $\sigma \approx 6$ km s $^{-1}$ is the line-of-sight velocity dispersion of the stars in the centre of the dwarf.

The solid line in the lower panel of Fig. 4 shows the rotation curve obtained from such a data set by binning the data in a way analogous to the one applied to the data for Leo I. Note that our simulated dwarf galaxy initially had a stellar disk which during the tidal evolution was transformed into a bar and then to a spheroid. In the final stage some residual rotation is still present as verified in Fig. 4. The equatorial plane of the dwarf (perpendicular to its total angular momentum vector) is inclined by about 60 degrees to the orbital plane so we do not actually see the maximum rotation.

Fig. 4 clearly demonstrates that a similar effect of reversed rotation as the one seen in Leo I can be produced by the presence of strong tidal tails. The intrinsic rotation well visible in the inner part of the simulated dwarf is reduced as we go towards larger $|x|$ and gets reversed at about ± 1.5 kpc which is well inside the dwarf. Note that the radius of the dwarf where the density profile starts to flatten due to tidal tails is about 2.5 kpc (Klimentowski et al. 2007) which corresponds to about 10 arcmin for Leo I. This behaviour is caused by the kinematics of stars in the tails: the stars typically move away from the dwarf with velocities proportional to their distance (see fig. 21 of Klimentowski et al. 2007). In the configuration presented in Fig. 4 this motion is in the opposite direction with respect to the intrinsic rotation of the dwarf on both sides of the galaxy. In the inner parts the stars tracing the intrinsic kinematics of the dwarf dominate and the rotation is well visible. When moving away from the centre there is a point where the stars from the tidal tails start to prevail and the dominant motion changes direction.

In choosing the stars to calculate the rotation curve we made only a simple constant cut-off in velocity with respect to the mean velocity of the dwarf. Although the sample of 328 stars in M08 was obtained in a similar way, in Fig. 3 we demonstrated that the behaviour of the rotation curve is preserved also for the sample of 316 stars cleaned with our method of interloper rejection. In order to check whether this is the case also for the simulated data we randomly selected 100 samples of 200 stars each from the total sample of stars used before and cleaned them of interlopers. The dashed line in Fig. 4 shows the average rotation curve calculated from these cleaned samples. As expected, the effect of the tidal tails is now less pronounced, but the reversed rotation is still present. We verified using the full 3D information from the simulation that the result shown with the dashed line would be almost identical if we actually removed all unbound stars. The reason for this is that some of the stars in the tails are still bound to the dwarf while moving away from it. On the other hand, if the tidal tails are cut off by considering only the stars within $|y| < 2.5$ kpc, the reversed rotation disappears, as shown by the dotted line in Fig. 4. This proves that indeed the tidal tails are responsible for the reversal of rotation.

Although in Fig. 4 the rotation is reversed at around 1.5 kpc (or 1.8 kpc for the cleaned sample) corresponding to 0.6 of the dwarf radius, in Leo I it seems to occur at 3 arcmin, i.e. at a much smaller fraction of radius equal to 0.3. Such differences are expected in light of the fact that the simulated dwarf was never intended to be a precise model of Leo I. It is simply employed to propose a plausible model for the kinematics of Leo I. The radius of rotation reversal could be easily changed by varying simulation parameters such as the initial concentration of the halo or the initial

disk scale length of the stars which will affect the effective tidal radius (Mayer et al. 2002).

If the overall qualitative picture presented here is correct then the eastern side of Leo I must be affected by the trailing tidal tail while the western side by the leading tail. Note that the rotation curves for 328 stars and 316 stars (left and right panels of Fig. 3) are quite similar. This further confirms our suspicion from the previous section that the sample of 316 stars is still to some extent contaminated by tidal tail stars. This contamination, however subtle, should still be taken into account; it supports our suggestion to include only the inner data points of the dispersion and kurtosis in the kinematic modelling of Leo I.

3.2 The symmetry of the velocity distribution

Another issue discussed by S07 and M08 is the question of the symmetry of the velocity distribution of the stellar sample of Leo I. While S07 found their distribution to be highly asymmetric, the one reported by M08 was quite symmetric and Gaussian-like. In the upper panel of Fig. 5 we show the distribution of velocities of stars along the y axis as measured by an observer situated in the same way with respect to the dwarf as before, now with the ± 30 km s⁻¹ cut-off in velocity, corresponding to 5σ range, larger this time to explore the tails of the velocity distribution. Although the distribution for the dwarf stars is embedded in a uniform background from the tails, it appears quite symmetric. In the middle panel we plot a similar distribution but now obtained from a sum of 100 samples of 200 stars each selected randomly from the previous one and cleaned of interlopers. On top of the distribution we plotted a Gaussian with dispersion of $\sigma = 5$ km s⁻¹ calculated from the sample. The lower panel shows the distribution of line-of-sight velocities of bound stars from the inside of the dwarf, i.e. with radii $r < 2.5$ kpc. Again, a Gaussian with $\sigma = 4.9$ km s⁻¹ calculated from the sample is plotted on top.

In all three cases, the distribution is highly symmetric which is understandable because in the present configuration the tidal tails contribute to both negative and positive velocities similarly. In addition, the two lower panels show remarkable similarity which means that our interloper removal scheme works adequately in removing obvious outliers (i.e. the uniform distribution of the stars from the tails). However, it should be kept in mind that the method does not remove all interlopers. Indeed, in a configuration similar to the one used here (observation along the tidal tails) and with the same initial cut-off in velocity, the scheme removes on average 80 percent of unbound stars (Klimentowski et al. 2007). The remaining contamination, although not apparent in overall distributions like the ones shown in Fig. 5, is still present and responsible for the reversed rotation as demonstrated by Fig. 4.

3.3 Tests of the method of velocity moments

We used the simulation data in the same configuration to test the reliability of the method used to model Leo I, based on fitting the velocity dispersion and kurtosis profiles. The method has been tested extensively in the context of modelling the kinematic samples of galaxy clusters and shown to




Figure 5. Upper panel: the distribution of velocities of 1.2×10^5 stars along the y axis in the same configuration as shown in Fig. 4. Middle panel: a similar distribution obtained from a sum of 100 samples of 200 stars selected randomly and cleaned of interlopers (in the end we have 1.6×10^4 stars). Lower panel: velocity distribution of 7.7×10^4 stars from the inside of the dwarf ($r < 2.5$ kpc). All histograms were normalized to unity. In the two lower panels the lines show the Gaussian distributions with dispersions measured from the data.

reproduce well the properties of simulated cluster-size dark matter haloes (Sanchis, Lokas & Mamon 2004; Lokas et al. 2006, 2007). Here we proceed in a similar way and generate 10 kinematic samples from the stellar component of our simulated dwarf of 330 stars each. The stars were selected randomly from the region corresponding to a projected radius of $R < 3$ kpc and with an initial cut-off in velocity of $\pm 17 \text{ km s}^{-1}$ with respect to the dwarf’s mean velocity, which corresponds to 3σ range, exactly as in the original data set of M08. An example of such data set is shown in the upper left panel of Fig. 6 for sample number 10 in a way analogous to the way we presented the data for Leo I in Fig. 1.

The data were binned with 30 stars per bin to obtain the profiles of the velocity moments also shown in Fig. 6. We then fitted the moments with the solutions of the Jeans

equations adopting the assumptions that mass follows light and anisotropy $\beta = \text{const.}$ The distribution of light was obtained by fitting the Sérsic profile to the projected distribution of the stars which gave $R_S = 0.54$ kpc and $m = 1$. The errors on the estimated parameters, the total mass and anisotropy, were read from probability contours analogous to the ones for Leo I shown in Fig. 2. The best-fitting parameters together with 1σ errors are presented in Fig. 7 for all samples, including sample 10. The horizontal solid lines in the four panels in the upper two rows of the Figure indicate the true values of the parameters measured from the full 3D information on the simulated dwarf: $M = 4.0 \times 10^7 M_\odot$ and $\beta = -0.13$ (see Klimentowski et al. 2007).

The left panels of Fig. 7 show results for the case when only the velocity dispersion profile is fitted. When the entire samples of 330 stars are considered (circles), the quality of the fit is generally poor. More specifically, compared to the true properties of the dwarf measured from the 3D simulation data (horizontal solid lines), the mass is overestimated and the anisotropy underestimated. In particular, in three cases out of ten, the best-fitting of β is $\beta < -1$, including the most discrepant case of sample 10 which has $\beta = -3.2$. A similar result was obtained for Leo I when the data were treated in the same way. When the data are cleaned of interlopers (we then use 300 stars in 10 bins) and again only the velocity dispersion profile is fitted (squares), the quality of the fit improves dramatically. The masses are now only slightly underestimated and anisotropies slightly overestimated which is due to the specific properties of the velocity distribution in the dwarf in this configuration (see Klimentowski et al. 2007).

The right panels of Fig. 7 show the corresponding results in the case when both velocity dispersion and kurtosis are fitted. As expected, the mass estimates are similar to those in the left panel. This is because it is the dispersion profile which is mainly sensitive to the mass. However, the situation is completely different for the anisotropy. Although for the entire samples (with interlopers) the quality of the fits is still poor, the best-fit anisotropy values are in close agreement with the true 3D values. Interestingly, analysis of both types of samples result in very similar values of anisotropy suggesting that contamination does not affect the determination of β when both dispersion and kurtosis are fitted. This is due to the fact that, contrary to common belief, the kurtosis is not much more affected by contamination than the dispersion. The reason for the similar effect of interlopers on kurtosis may be that it is constructed via dividing the fourth velocity moment by the fourth power of dispersion so the influence of contamination may partially cancel out. The contamination increases the measured values of kurtosis (see the lower panels of Fig. 6), but the dispersion and the kurtosis depend on the velocity anisotropy in a different way, so that more strongly increasing dispersion corresponds to more tangential orbits while more strongly increasing kurtosis corresponds to more radial orbits (see fig. 4 in Lokas et al. 2005). The kurtosis values inflated by interlopers thus lead to the situation when more radial orbits are preferred which balances the preference for tangential orbits due to the inflated dispersion profile.

In the case when both moments are fitted for samples cleaned of interlopers the overall quality of the fits is good and in eight out of ten cases the true values of the parameters




Figure 6. Upper panels: the kinematic data sets for sample 10 generated from the simulated dwarf. The left panel shows the original sample of 330 stars selected by the cut-off in velocity of $\pm 17 \text{ km s}^{-1}$ with respect to the dwarf's mean velocity, while the right one presents the sample of 312 stars obtained by rejection of interlopers (open circles). The middle (lower) panels show the velocity dispersion (kurtosis) profiles obtained from the corresponding samples with 30 stars per bin. The dashed lines show the best-fitting dispersion profiles when only the dispersion is fitted, while the solid lines plot the best-fitting profiles of the moments when the dispersion and kurtosis are fitted simultaneously.

are within the 1σ errors of the best-fitting parameters. We can thus be quite confident that the error estimates for Leo I reflect the real uncertainty in the parameters.

4 DISCUSSION

We have demonstrated that after the application of the interloper removal scheme and the inclusion of the fourth velocity moment the kinematic data for Leo I dSph galaxy can be reliably modelled using the solutions of the Jeans equations and its properties can be recovered. We find that the data are consistent with a simple model in which mass follows light and stellar orbits are close to isotropic with no need for an extended dark matter halo. The picture of Leo I emerging from our analysis is fully consistent with the tidal stirring scenario for the formation of dSph galax-

ies (Mayer et al. 2001). The fact that Leo I has been significantly tidally stirred and that fairly prominent tails are present is not in contradiction with the fact that it is still a gravitationally bound object nearly in equilibrium and has a significant dark matter halo. This dark matter halo is not extended, but rather truncated just outside the stellar component. The scenario predicts that extended dark haloes are removed after one or two tidal shocks during pericentre passages if only the pericentre is small enough, i.e. below 50 kpc (Mayer et al. 2001, 2002). Also the intrinsic rotation present in Leo I is naturally explained within this scenario as a remnant of the initial disk.

The study of the rotation curve in Leo I shows that the cleaned sample is still contaminated to some extent by tidal tails and therefore it is advisable to use only the kinematic data from the inner part of the dwarf for dynamical modelling. The rotation curve constructed from the data, both




Figure 7. Results of fitting the velocity moments for 10 mock data samples generated from the simulated dwarf. The left panels show the best-fitting mass (upper panel) and anisotropy (middle panel) with 1σ errors and the goodness of fit measure (lower panel) in the case when only the dispersion profile is fitted. The right panels show the corresponding results in the case when both moments are fitted. The circles correspond to the results obtained for the entire samples, while the squares to those for samples cleaned of interlopers. Horizontal solid lines indicate the true values of the parameters measured from the 3D information.

for the initial and cleaned samples, shows that the rotation is reversed when going from the inside to the outside of the galaxy. We interpret this behaviour as due to the presence of contamination from the leading tidal tail in the western part of Leo I (seen in the background from the point of view of the observer) and from the trailing tail in the eastern part of the galaxy (seen in the foreground of the observer).

This kind of behaviour in the rotation curve may also reflect the presence of a counterrotating core. Such cores are however typically found in much brighter elliptical galaxies, an example is the galaxy NGC 770 studied in detail by Geha, Guhathakurta & van der Marel (2005). The rotation curve of the galaxy, shown in their fig. 3 looks very similar to the rotation curve of Leo I in Fig. 3 of the present paper and the photometric analysis shows that it is generated by the presence of a small inner disk. For the inner rotation of the disk to be well visible it has to be viewed close to edge-on. Then the inner contours of the surface brightness should be

more disky than the outer ones, i.e. the ellipticity should decrease with radius. It is indeed the case for NGC 770, as demonstrated by fig. 7 of Geha et al. (2005), but not for Leo I: as shown in fig. 17 of M08 the ellipticity of Leo I increases with radius, i.e. the inner contours of the surface density of the stars are more circular. This could only be reconciled with counterrotation if there is a counterrotating bar in Leo I viewed along the long axis. It is unclear how such a bar could form in the standard tidal stirring scenario for the formation of dSphs (Mayer et al. 2001). Although in the simulation employed here the dwarf has a bar for most of the time (it is destroyed only at the last pericentre, see Klimontowski et al. 2008), it rotates in the same direction as the rest of the stars. One possibility of creating a counterrotating bar is through an interaction of Leo I with some other dwarf galaxy in the past (e.g. Kravtsov, Gnedin & Klypin 2004).

In light of the analysis presented here, Leo I looks very similar to the Fornax dwarf for which it is also found that




Figure 8. Velocity dispersion profiles expected for the samples of stars with different measurement errors. The solid lines correspond to the exact measurements, the dashed lines to the measurements with Gaussian errors of 2 km s^{-1} and the dotted lines to the errors of 5 km s^{-1} . The left panel is for the entire sample of 328 stars, the right panel shows analogous results after removal of interlopers.

after the removal of interlopers the model with mass following light and stellar orbits close to isotropic provides satisfactory description of the kinematics (Klimontowski et al. 2007; Lokas, Klimontowski & Wojtak 2007). Both dwarfs also have a rather low mass-to-light ratio compared to systems like Draco. The important difference is the source of contamination in the kinematic samples: while in Fornax the majority of contamination probably comes from Milky Way stars, in Leo I it is due to tidal tails. In the direction of Leo I the contamination from the Milky Way is negligible. According to the Besancon model of the Milky Way (Robin et al. 2003), the stars from our Galaxy are expected to have heliocentric velocities below 100 km s^{-1} . This corresponds roughly to a 200 km s^{-1} difference with respect to the mean velocity of Leo I stars.

Our estimates of the mass and mass-to-light ratio agree within errors with those of M08 and S07 but are by a factor of a few lower than those of K07. Once the difference in the (much lower) assumed luminosity of K07 is taken into account, the discrepancy is alleviated (K07 assumed $L_V = 3.4 \times 10^6 M_\odot$ which should have been corrected for the much lower distance adopted by Irwin & Hatzidimitriou 1995). We fitted the dispersion profile from the lower right panel of fig. 8 in K07 with our adopted parameters (see Table 1) and assumptions (mass follows light and $\beta = \text{const}$). We find that our model fits their data well (although with a very tangential anisotropy $\beta = -4.4$) for a value of mass as low as $5.0 \times 10^7 M_\odot$ which corresponds to $M/L_V = 9.1 M_\odot/L_\odot$. This value differs by ~ 40 percent from our preferred value of $M/L_V = 6.6$.

Comparing our velocity dispersion profile from M08 (even for the original contaminated sample) to that of K07, we find that the latter is typically higher (note however that, contrary to the data sets of M08 and S07, no secondary increase of the dispersion profile at larger radii is seen). This may be due to errors in velocity measurements which are of the order of 5 km s^{-1} in K07, compared to 2 km s^{-1} in M08. To verify this possibility we calculated the dispersion profiles again from the M08 sample modifying the velocities by errors generated from a Gaussian distribution with a dispersion of 2 and 5 km s^{-1} . For each error value a hundred samples were generated from which the dispersions were calculated and averaged. The results are shown in the left panel of Fig. 8. The right panel of the Figure shows analogous re-

sults except that the procedure of interloper removal was applied before calculating the dispersions and averaging. It is clear, that while the errors of 2 km s^{-1} have negligible impact, those of 5 km s^{-1} increase the velocity dispersion by 1.2 km s^{-1} on average. Decreasing the dispersion values in K07 by this correction yields the best fitting mass of $3.8 \times 10^7 M_\odot$ which corresponds to $M/L_V = 6.8 M_\odot/L_\odot$ in excellent agreement with our results in Table 2. The larger mass-to-light ratio inferred by K07 may therefore be due to adopting lower luminosity and larger velocity errors, and considering models with extended dark matter haloes.

Using an N -body simulation we have shown that the line-of-sight velocity distribution of a dwarf near apocentre should be symmetric and close to Gaussian in shape, as is indeed the case for the stellar sample of M08 for Leo I. The reason why S07 found the distribution of their sample to be asymmetric probably lies in their non-uniform coverage of the galaxy or the bias in velocity measurements. M08 compared their velocity measurements to those of S07 for the stars common to both samples and found (see their fig. 7) a velocity offset dependent on the heliocentric velocity such that in S07 the velocities from below the galaxy mean velocity are shifted towards the mean (even by 10 km s^{-1} for heliocentric velocities of 270 km s^{-1}). This may be the reason why the low-velocity tail of the distribution disappears and the resulting distribution is not symmetric (see fig. 14 of S07).

Departures from symmetry in the velocity distribution are expected at large distances from the centre due to the asymmetry of tidal tails. However, as pointed out by M08, any contamination from tidal tails in Leo I probably comes from regions close to the dwarf, where the tails and the velocity distribution are symmetric. This is because the usual features seen in the colour-magnitude diagram of Leo I are well visible (they would be blurred if the stars came from a wide variety of distances). When studying the velocity distribution in the N -body simulation, we introduced a velocity cut-off of $\pm 30 \text{ km s}^{-1}$ which corresponds to distances of tidal tail stars less than 20 kpc (see fig. 21 in Klimontowski et al. 2007). Based on photometric measurements, M08 estimate the tidal extensions of Leo I to correspond to distances less than 40 kpc , which means that our choice is fairly conservative and consistent with their photometric constraints.

The above discussion provides further support for the

symmetry of the velocity distribution in Leo I. Indeed, S07 considered the velocity distribution of tidal tails formed in their N -body models out to much larger distances and found that the trailing arm has a more extended velocity tail than the leading one. Comparing the velocity distributions of their N -body models with that of Leo I, S07 concluded that the latter must move to the east, i.e. the leading tidal tail is on the eastern side of the dwarf. In this case, the reversed rotation in Fig. 3 cannot be explained (the velocities in the leading arm would have to be towards the observer so Leo I would have to be approaching instead of receding). If indeed the biased velocity measurements are responsible for the asymmetry of the velocity distribution in the study of S07 then the distribution may in fact be symmetric and no information on the orbit and the orientation of the tidal tails can be inferred from it alone.

ACKNOWLEDGEMENTS

We are grateful to M. Mateo et al. for providing the kinematic data for Leo I stars in electronic form. SK is funded by the Center for Cosmology and Astro-Particle Physics at The Ohio State University. The numerical simulations were performed on the zBox1 supercomputer at the University of Zürich. We made use of the Besancon Galaxy model available at <http://bison.obs-besancon.fr/modele/>. This research was partially supported by the Polish Ministry of Science and Higher Education under grant N N203 0253 33.

REFERENCES

- Aaronson M., Hodge P. W., Olszewski E. W., 1983, *ApJ*, 267, 271
 Bellazzini M., Gennari N., Ferraro F. R., Sollima A., 2004, *MNRAS*, 354, 708
 Byrd G., Valtonen M., McCall M., Innanen K., 1994, *AJ*, 107, 2055
 Caputo F., Cassisi S., Castellani M., Marconi G., Santolamazza P., 1999, *AJ*, 117, 2199
 den Hartog R., Katgert P., 1996, *MNRAS*, 279, 349
 Gallart C., Freedman W. L., Aparicio A., Bertelli G., Chiosi C., 1999, *AJ*, 118, 2245
 Geha M., Guhathakurta P., van der Marel R. P., 2005, *AJ*, 129, 2617
 Harrington R. G., Wilson A. G., 1950, *PASP*, 62, 118
 Held E. V., Clementini G., Rizzi L., Momany Y., Saviane I., Di Fabrizio L., 2001, *ApJ*, 562, L39
 Irwin M., Hatzidimitriou D., 1995, *MNRAS*, 277, 1354
 Klimentowski J., Lokas E. L., Kazantzidis S., Prada F., Mayer L., Mamon G. A., 2007, *MNRAS*, 378, 353
 Klimentowski J., Lokas E. L., Kazantzidis S., Mayer L., Mamon G. A., Prada F., 2008, submitted to *MNRAS*, arXiv:0803.2464
 Koch A., Wilkinson M. I., Kleyna J. T., Gilmore G. F., Grebel E. K., Mackey A. D., Evans N. W., Wyse R. F. G., 2007, *ApJ*, 657, 241
 Kravtsov A. V., Gnedin O. Y., Klypin A. A., 2004, *ApJ*, 609, 482
 Lee M. G., Freedman W., Mateo M., Thompson I., Roth M., Ruiz M. T., 1993, *AJ*, 106, 1420
 Lokas E. L., 2002, *MNRAS*, 333, 697
 Lokas E. L., Mamon G. A., 2003, *MNRAS*, 343, 401
 Lokas E. L., Mamon G. A., Prada F., 2005, *MNRAS*, 363, 918
 Lokas E. L., Wojtak R., Gottlöber S., Mamon G. A., Prada F., 2006, *MNRAS*, 367, 1463
 Lokas E. L., Wojtak R., Mamon G. A., Gottlöber S., 2007, arXiv:0712.2368
 Lokas E. L., Klimentowski J., Wojtak R., 2007, arXiv:0712.2372
 Mateo M., Olszewski E. W., Vogt S. S., Keane M. J., 1998, *AJ*, 116, 2315
 Mateo M., Olszewski E. W., Walker M. G., 2008, *ApJ*, 675, 201
 Mayer L., Governato F., Colpi M., Moore B., Quinn T., Wadsley J., Stadel J., Lake G., 2001, *ApJ*, 559, 754
 Mayer L., Moore B., Quinn T., Governato F., Stadel J., 2002, *MNRAS*, 336, 119
 Robin A. C., Reylé C., Derrière S., Picaud S., 2003, *A&A*, 409, 523
 Sales L. V., Navarro J. F., Abadi M. G., Steinmetz M., 2007, *MNRAS*, 379, 1475
 Sanchis T., Lokas E. L., Mamon G. A., 2004, *MNRAS*, 347, 1198
 Sérsic J. L., 1968, *Atlas de Galaxies Australes*, Observatorio Astronómico, Córdoba
 Sohn S. T. et al., 2007, *ApJ*, 663, 960
 Suntzeff N. B., Aaronson M., Olszewski E. W., Cook K. H., 1986, *AJ*, 91, 1091